

Gender Differences and Stimulant Use Supplemental Module Reference List

- American College of Obstetricians and Gynecologists, Committee Opinion No. 479: Methamphetamine abuse in women of reproductive age. (2011). *Obstetrics & Gynecology*, 117(3), 751–755. doi: 10.1097/aog.0b013e318214784e
- Anker, J. J., & Carroll, M. E. (2010). Females are more vulnerable to drug abuse than males: Evidence from preclinical studies and the role of ovarian hormones. *Biological Basis of Sex Differences in Psychopharmacology Current Topics in Behavioral Neurosciences*, 73–96. doi: 10.1007/7854_2010_93
- Anthony, J. C., Warner, L. A., & Kessler, R. C. (1994). Comparative epidemiology of dependence on tobacco, alcohol, controlled substances, and inhalants: Basic findings from the National Comorbidity Survey. *Experimental and Clinical Psychopharmacology*, 2(3), 244–268. doi: 10.1037/1064-1297.2.3.244
- Bauer, C. R., Langer, J. C., Shankaran, S., Bada, H. S., Lester, B., Wright, L. L., ... Verter, J. (2005). Acute neonatal effects of cocaine exposure during pregnancy. *Archives of Pediatrics & Adolescent Medicine*, 159(9), 824. doi: 10.1001/archpedi.159.9.824
- Beck, J.S. (2011). *Cognitive Behavior Therapy*. Basics and beyond (2nd ed.). New York, NY: Guilford Press.
- Betancourt, L. M., Yang, W., Brodsky, N. L., Gallagher, P. R., Malmud, E. K., Giannetta, J. M., ... Hurt, H. (2011). Adolescents with and without gestational cocaine exposure: Longitudinal analysis of inhibitory control, memory and receptive language. *Neurotoxicology and Teratology*, 33(1), 36–46. doi: 10.1016/j.ntt.2010.08.004
- Brecht, M.-L., O'Brien, A., Mayrhauser, C. V., & Anglin, M. (2004). Methamphetamine use behaviors and gender differences. *Addictive Behaviors*, 29(1), 89–106. doi: 10.1016/s0306-4603(03)00082-0
- Brizendine, L. (2009). *The female brain*. London: Bantam.
- Cain, M. A., Bornick, P., & Whiteman, V. (2013). The maternal, fetal, and neonatal effects of cocaine exposure in pregnancy. *Clinical Obstetrics and Gynecology*, 56(1), 124–132. doi: 10.1097/grf.0b013e31827ae167
- Campbell, U. C., Morgan, A. D., & Carroll, M. E. (2002). Sex differences in the effects of baclofen on the acquisition of intravenous cocaine self-administration in rats. *Drug and Alcohol Dependence*, 66(1), 61–69. doi: 10.1016/s0376-8716(01)00185-5
- Cretzmeyer, M., Sarrazin, M. V., Huber, D. L., Block, R. I., & Hall, J. A. (2003). Treatment of methamphetamine abuse: Research findings and clinical directions. *Journal of Substance Abuse Treatment*, 24(3), 267–277. doi: 10.1016/s0740-5472(03)00028-x
- Diaz, S. D., Smith, L. M., Lagasse, L. L., Derauf, C., Newman, E., Shah, R., ... Lester, B. M. (2014). Effects of prenatal methamphetamine exposure on behavioral and cognitive findings at 7.5 years of age. *The Journal of Pediatrics*, 164(6), 1333–1338. doi: 10.1016/j.jpeds.2014.01.053
- Dluzen, D. E., & Liu, B. (2008). Gender differences in methamphetamine use and responses: A review. *Gender Medicine*, 5(1), 24–35. doi: 10.1016/s1550-8579(08)80005-8

- Evans, S. M., & Foltin, R. W. (2005). Exogenous progesterone attenuates the subjective effects of smoked cocaine in women, but not in men. *Neuropsychopharmacology*, *31*(3), 659–674. doi: 10.1038/sj.npp.1300887
- Eze, N., Smith, L. M., Lagasse, L.L., Derauf, C., Newman, E., Arria, A., ... Lester, B. (2016). School-aged outcomes following prenatal methamphetamine exposure: 7.5-year follow-up from the Infant Development, Environment, and Lifestyle Study. *Journal of Pediatrics*, *170*, 34–38. doi: 10.1016/j.jpeds.2015.11.070
- Eyler, F. D., Behnke, M., Garvan, C. W., Woods, N. S., Wobie, K., & Conlon, M. (2001). Newborn evaluations of toxicity and withdrawal related to prenatal cocaine exposure. *Neurotoxicology and Teratology*, *23*(5), 399–411. doi: 10.1016/s0892-0362(01)00166-0
- Gorman, M. C., Orme, K. S., Nguyen, N. T., Kent, E. J., & Caughey, A. B. (2014). Outcomes in pregnancies complicated by methamphetamine use. *American Journal of Obstetrics and Gynecology*, *211*(4), 429 e.1–429.e7. doi: 10.1016/j.ajog.2014.06.005
- Hachtel, J. C., & Armstrong, K. J. (2019). Illicit use of prescription stimulants: Gender differences in perceptions of risk. *Substance Use & Misuse*, *54*(10), 1654-1662. doi: 10.1080/10826084.2019.1608246
- Jackson, L. R., Robinson, T. E., & Becker, J. B. (2005). Sex differences and hormonal influences on acquisition of cocaine self-administration in rats. *Neuropsychopharmacology*, *31*(1), 129–138. doi: 10.1038/sj.npp.1300778
- Kampman, K. (2008). The search for medications to treat stimulant dependence. *Addiction Science & Clinical Practice*, *4*(2), 28–35. doi: 10.1151/ascp084228
- Kippin, T. E., Fuchs, R. A., Mehta, R. H., Case, J. M., Parker, M. P., Bimonte-Nelson, H. A., & See, R. E. (2005). Potentiation of cocaine-primed reinstatement of drug seeking in female rats during estrus. *Psychopharmacology*, *182*(2), 245–252. doi: 10.1007/s00213-005-0071-y
- Lagasse, L. L., Derauf, C., Smith, L. M., Newman, E., Shah, R., Neal, C., ... Lester, B. M. (2012). Prenatal methamphetamine exposure and childhood behavior problems at 3 and 5 years of age. *Pediatrics*, *129*(4), 681-688. doi: 10.1542/peds.2011-2209
- Maxwell, J. C. (Personal Communication October 3, 2019). Reanalysis of methamphetamine survey data.
- Maxwell, J. C. (2014). A new survey of methamphetamine users in treatment: Who they are, why they like “meth,” and why they need additional services. *Substance Use & Misuse*, *49*(6), 639–644. doi: 10.3109/10826084.2013.841244
- National Institute on Drug Abuse. (2016). *Cocaine*. Retrieved from <https://www.drugabuse.gov/drugs-abuse/cocaine> on 2020, May 15.
- National Institute on Drug Abuse. (2000). *Gender Differences in Drug Abuse Risks and Treatment*. Retrieved from <https://archives.drugabuse.gov/news-events/nida-notes/2000/09/gender-differences-in-drug-abuse-risks-treatment> on 2020, May 1.

Rawson, R. A., Shoptaw, S. J., Obert, J. L., Mccann, M. J., Hasson, A. L., Marinelli-Casey, P. J., ... Ling, W. (1995). An intensive outpatient approach for cocaine abuse treatment. *Journal of Substance Abuse Treatment*, 12(2), 117–127. doi: 10.1016/0740-5472(94)00080-b

Robbins, S. J., Ehrman, R. N., Childress, A. R., & O'Brien, C. P. (1999). Comparing levels of cocaine cue reactivity in male and female outpatients. *Drug and Alcohol Dependence*, 53(3), 223–230. doi: 10.1016/s0376-8716(98)00135-5

Rowan-Szal, G. A., Joe, G. W., Simpson, D. D., Greener, J. M., & Vance, J. (2009). During-treatment outcomes among female methamphetamine-using offenders in prison-based treatments. *Journal of Offender Rehabilitation*, 48(5), 388–401. doi: 10.1080/10509670902979496

Smith, L. M., Lagasse, L. L., Derauf, C., Grant, P., Shah, R., Arria, A., ... Lester, B. M. (2006). The Infant Development, Environment, and Lifestyle Study: Effects of prenatal methamphetamine exposure, polydrug exposure, and poverty on intrauterine growth. *Pediatrics*, 118(3), 1149–1156. doi: 10.1542/peds.2005-2564

Smith, T. E., Desantis, A. D., & Martel, M. M. (2017). Gender differences in nonprescribed psychostimulant use in young adults. *Substance Use & Misuse*, 53(4), 622–628. doi: 10.1080/10826084.2017.1355384

Substance Abuse and Mental Health Services Administration. (2019). *Key Substance Use and Mental Health Indicators in the United States: Results from the 2018 National Survey on Drug Use and Health* (HHS Publication No. PEP19-5068, NSDUH Series H-54). Rockville, MD: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration. Retrieved from <https://www.samhsa.gov/data/report/2018-nsduh-annual-national-report> and [https://www.samhsa.gov/data/sites/default/files/reports/rpt23250/5 Women 2020 01 14 508.pdf](https://www.samhsa.gov/data/sites/default/files/reports/rpt23250/5_Women_2020_01_14_508.pdf) on 2020, May 15.

Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality. (2017). *Results from the 2016 National Survey on Drug Use and Health: Detailed Tables*. Rockville, MD: Substance Abuse and Mental Health Services Administration. Retrieved from <https://www.samhsa.gov/data/sites/default/files/NSDUH-DetTabs-2016/NSDUH-DetTabs-2016.pdf> on 2020, May 15.

Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality. (2014). *The TEDS Report: Gender Differences in Primary Substance of Abuse across Age Groups*. Rockville, MD: Substance Abuse and Mental Health Services Administration. Retrieved from https://www.dasis.samhsa.gov/dasis2/teds_pubs/2014_teds_rpt_natl.pdf on 2020, May 15.

Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality. (2016). *Treatment Episode Data Set (TEDS): 2004-2014. National Admissions to Substance Abuse Treatment Services*. Rockville, MD: Substance Abuse and Mental Health Services Administration. Retrieved from https://www.dasis.samhsa.gov/dasis2/teds_pubs/2014_teds_rpt_natl.pdf on 2020, May 15.

Volkow, N. (April 21, 2005). Testimony before the Subcommittee on Labor, Health, and Human Services; Education; and Related Agencies. Committee on Appropriations, U.S. Senate.

Zweben, J. E., Cohen, J. B., Christian, D., Galloway, G. P., Salinardi, M., Parent, D., & Iguchi, M. (2004). Psychiatric symptoms in methamphetamine users. *American Journal on Addictions*, *13*(2), 181–190. doi: 10.1080/10550490490436055