



TRANSITIONAL AGE YOUTH (TAY)
webinar series

Who's Doing What?: The Epidemiology of Adolescent Substance Use

Produced in Partnership by:



Network Coordinating Office

ATTC

Addiction Technology Transfer Center Network
Funded by Substance Abuse and Mental Health Services Administration



Interdisciplinary Leaders in
Substance Use Education,
Research, Care and Policy

ADOLESCENT

S B I R T

Screening, Brief Intervention & Referral to Treatment

by

NORC

at the
University of
Chicago

Webinar Moderator

Tracy McPherson, PhD

Senior Research Scientist


Public Health Department

NORC at the University of Chicago

McPherson-Tracy@norc.org



 **NORC** at the
University of
Chicago



Produced in
Partnership...



Network Coordinating Office

ATTC

Addiction Technology Transfer Center Network
Funded by Substance Abuse and Mental Health Services Administration



Interdisciplinary Leaders in
Substance Use Education,
Research, Care and Policy


ADOLESCENT
SBIRT

Screening, Brief Intervention & Referral to Treatment

by

NORC

at the
University of
Chicago





TRANSITIONAL AGE YOUTH (TAY)

webinar series


- 1) The Impact of Substance Use on the Developing Adolescent Brain
- 2) Who's Doing What?: The Epidemiology of Adolescent Substance Use
- 3) Substance Use Interventions for Adolescents and Transitional Age Youth
- 4) Integrating Loss and Grief into the SBIRT Model

attcnetwork.org/centers/global-attc/tay-webinar-series

Access Materials

- Bookmark the website
- Follow-up email
- On-demand access 24/7
- Brief survey
- Certificate of Completion brief application (1 NAADAC CE)



sbirt.webs.com/adolescent-epidemiology



Access PowerPoint Slides

Access the PowerPoint slides through the “Handouts” pane of your GoToWebinar Control Panel on your computer or mobile device.



Or, download them from the website.



Ask Questions

Ask questions at any point through the “Chat” pane of your GoToWebinar Control Panel on your computer or mobile device.

Answers will be posted on the website.



Webinar Presenter


Veronika Mesheriakova, MD

Assistant Professor

Pediatrics and Adolescent Medicine,
University of California, San Francisco

Director

UCSF Youth Outpatient Substance Use
Program (YoSUP)



Objectives


After completing this activity, learners will be able to:

- Discuss trends in identification and treatment of adolescent substance use
- Describe the implications of adolescent substance use
- Outline substances most commonly used by adolescents in the U.S. and describe trends in adolescent substance use over time
- Relate how the family and positive social supports serve as protective factors.



A Case

- Breanne is a 16yo young woman who moved to a new town with her family and presented to establish care with a local pediatrician.
- During the initial visit, she complained of insomnia, fatigue, loss of appetite (with a 20lb weight loss), she reported that she quit her volleyball team, and has been failing her classes.
- She reported that she had been using cannabis to help with insomnia.
- She was diagnosed with a depressive disorder and started on fluoxetine at that time.




Case Continued

- Two weeks later, Breanne saw her pediatrician again and reported doing a bit better in school, was less irritable, and was sleeping a bit better.
- She was also excited about a family trip to Europe but expressed significant anxiety about flying.
- Her pediatrician discussed non-pharmacologic options and prescribed alprazolam for flight anxiety and counseled parents to keep the medication locked, and in their possession at all times.


Case Continued

- Two months later, Breanne had another office visit. She had a great time on vacation and reported that her mood was much better.
- She stated that she hadn't used cannabis for a few weeks because she couldn't buy it while on vacation.



Case Continued


- Six months later, Breanne continued to struggle in school and started coming home visibly high. She eventually admitted to her parents that she had been using opioids that she purchased from friends.
- During private interview, Breanne reported a 4 year history of substance use that started with cannabis but later came to include nicotine, alcohol, cocaine, LSD, and various prescription medications.
- She was eventually diagnosed with:
 - Opioid use disorder
 - Cannabis use disorder
 - Benzodiazepine use disorder
 - Stimulant use disorder






Detection and Treatment of Youth SUDs

- Only about 45% of pediatricians routinely screen adolescents for substance use
- Only 16% of pediatricians screen using a validated screening tool
- Pediatricians tend to underestimate the severity of adolescents' substance use problems
 - One study revealed that only 4.8% of patients with problematic substance use were correctly identified
- Even when problem substance use is identified, up to 20% of those patients receive no intervention



Medical Education on SUDs: Medical School


- On average, medical school curricula offer a total of 12 hours to education on substance use and addiction
 - Most of this training is related to scientific pathophysiology and very little is focused on attitudes/stigma, screening, or evidence-based treatment of addiction
- Only 12 medical schools in the U.S. require a course on addiction
- Only 45 medical schools in the U.S. offer an elective course on addiction
- Medical students' attitudes about people with addiction tend to become increasingly negative as they progress through training.




Medical Education on SUDs: Residency

Pediatrics ACGME requirements:

- ...instruction and experience in pain management if applicable for the specialty, including recognition of the signs of addiction




Substance Use and Addiction is a Pediatric Priority



A Pediatric Priority: Burden of Disease


- Worldwide, mental and substance use disorders are the **#1** contributors to health burden among 10-24 year-olds (19% of all disease burden)
- Among young adults aged 20-24, alcohol use is the largest worldwide contributor to disease burden (10% of all disease burden).
- In the U.S. almost $\frac{1}{4}$ of all deaths are attributable to (directly or indirectly) to the use of substances.



A Pediatric Priority: Adolescents' Vulnerability

- Adolescence is a neurobiologically vulnerable time period for the development of substance use disorders
 - Fully developed pain and reward pathways
 - Immature prefrontal cortex

Images of Brain Development in Healthy Children and Teens (Ages 5-20)




The brain continues to develop through early adulthood. Mature brain regions at each developmental stage are indicated in blue. The prefrontal cortex (red circles), which governs judgment and self-control, is the last part of the brain to mature.

Source: *PNAS* 101:8174-8179, 2004.

A Pediatric Priority: Adolescents' Vulnerability


Adolescents are the group that is most likely to experience health consequences related to substance use

- Accidents
- Dating violence
- Risky sexual practices
- Development of substance use disorders



Addiction as a Pediatric-Onset Illness


- 90% adults who suffer from a substance use disorder started using substances before age 18 and developed their substance use disorder before age 20
 - 50% started using substances before age 15
- Younger age at first substance use is a strong risk factor for the development of substance use disorders
 - 15.4% of people who had their first drink before age 14 will develop a use disorder vs. only 2.1% of people who had their first drink after age 21
 - 25% of people who first misuse Rx drugs before age 13




Addiction as a Pediatric-Onset Illness

Predictors of illness duration

- Age of first use
 - Started using after age 20 → 18 years of illness
 - Started using before age 15 → 29 years of illness
- Duration of use before first treatment
 - 10 years of use before first treatment → 15 years of illness
 - 20 years of use before first treatment → 35+ years of illness



Epidemiology of Substance Use in the United States



Poll

How many people aged 12 and older in the United States reported using substances in the past month (tobacco, alcohol, other drugs)?


A. 165 million (60%)

B. 140 million (51%)

C. 58 million (21%)

D. 36 million (13%)

Figure 1. Past Month Substance Use among People Aged 12 or Older: 2019



Rx = prescription.

Note: Substance Use includes any illicit drug, kratom, alcohol, and tobacco use.

Note: The estimated numbers of current users of different substances are not mutually exclusive because people could have used more than one type of substance in the past month.

Poll

How many people aged 12 and older in the United States have had a substance use disorder in the past year?

- A. 5 million
- B. 10 million
- C. 15 million
- D. 20 million





Figure 46. People Aged 12 or Older with a Past Year Substance Use Disorder (SUD): 2019



Note: The estimated numbers of people with substance use disorders are not mutually exclusive because people could have use disorders for more than one substance.

Addiction as an Intergenerational Issue

- 1 in 8 of all U.S. children are living in a household where at least one parent has struggled with a substance use disorder (SUD) in the past year.






Figure 2. Number and percentage of children aged 17 or younger living with at least one parent with a past year alcohol use disorder, by age group and household composition: annual average, 2009 to 2014



Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Surveys on Drug Use and Health (NSDUHs), 2009 to 2014.


Figure 3. Number and percentage of children aged 17 or younger living with at least one parent with a past year illicit drug use disorder, by age group and household composition: annual average, 2009 to 2014




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Surveys on Drug Use and Health (NSDUHs), 2009 to 2014.


Addiction as an Intergenerational Issue

- Children with parents struggling with SUDs are more likely to:
 - Have low socio-economic status
 - Struggle more with academic, social, and family functioning
 - Experience mental and behavioral disorders
 - Develop SUDs themselves
 - Children of parents with alcohol use disorder (AUD) are 4x more likely to develop AUD than their peers who don't have a parent with AUD
 - Parental marijuana use increases risk of adolescent marijuana, tobacco, and alcohol use and opioid misuse



Epidemiology of Adolescent Substance Use





Monitoring the Future is an annual survey of 8th, 10th, and 12th graders conducted by researchers at the Institute for Social Research at the University of Michigan, Ann Arbor, under a grant from the National Institute on Drug Abuse, part of the National Institutes of Health. Since 1975, the survey has measured how teens report their drug, alcohol, and cigarette use and related attitudes in 12th graders nationwide; 8th and 10th graders were added to the survey in 1991.

42,531 STUDENTS FROM 396 PUBLIC AND PRIVATE SCHOOLS PARTICIPATED IN THE 2019 SURVEY.

Poll


Which substance is most commonly used by adolescents

A. Nicotine

B. Alcohol

C. Cannabis


D. Prescription drugs



Poll

What has been the trend in rates of alcohol use among 12th graders over the past 10 years?


- A. Rates have decreased
- B. Rates have increased
- C. Rates have stayed about the same



ALCOHOL USE CONTINUES ITS DECLINE


PAST YEAR ALCOHOL USE

Significant long-term decrease in all grades



BINGE DRINKING*

Significant long-term decrease in all grades



8th graders

10th graders

12th graders

*5 or more drinks in a row in the past two weeks



DRUGABUSE.GOV


Poll

What percentage of 12th graders reported vaping nicotine on a daily basis?

A. Almost 7%


B. Almost 12%

C. Almost 20%



TOBACCO AND NICOTINE: VAPING THREATENS PROGRESS

NICOTINE – DAILY USE



*Significant decline from 2018 (3.6%)

CIGARETTE SMOKING (PAST MONTH) DECLINES OVER PAST TEN YEARS



*Significant decline from 2018 (7.6%)

TO VIEW MORE RESULTS ON VAPING VISIT:

<https://www.drugabuse.gov/related-topics/trends-statistics/infographics/monitoring-future-2019-survey-results-vaping>




DRUGABUSE.GOV

Poll

What has been the trend in rates of illicit substance use (including cannabis) among 12th graders over the past 20?


- A. Rates have decreased
- B. Rates have increased
- C. Rates have stayed about the same



ILLICIT DRUG USE


ILLICIT DRUG USE STEADY

Past year use among 12th graders



PAST YEAR ILLICIT DRUG USE

Past year use among 12th graders




National Institute on Drug Abuse

DRUGABUSE.GOV

Poll

What has been the trend in rates of daily cannabis use among adolescents since 2017?


- A. Rates have increased in younger adolescents
- B. Rates have decreased in older adolescents
- C. Rates have stayed about the same in younger adolescents



DAILY MARIJUANA USE IN LOWER GRADES INCREASES BUT PAST YEAR MARIJUANA USE STEADY


DAILY MARIJUANA USE

sees significant increase among 8th and 10th graders since 2018



PAST YEAR MARIJUANA USE

gap closing between older grades



8th graders 10th graders 12th graders




DRUGABUSE.GOV



Poll

What has been the trend in prescription drug misuse among adolescents over the past 10 years?

- A. Rates have increased
- B. Rates have decreased
- C. Rates have stayed about the same



PRESCRIPTION DRUG MISUSE CONTINUES DECLINE FROM PEAK YEARS



8th graders 10th graders 12th graders




DRUGABUSE.GOV

PRESCRIPTION DRUG MISUSE CONTINUES DECLINE FROM PEAK YEARS

ADDERALL MISUSE SEES SIGNIFICANT CHANGES IN PAST 5 YEARS

a decrease in 10th and 12th grades, but an increase in 8th grade



DRUGABUSE.GOV

Substance Use Disorders Among Adolescents






Figure 45. Substance Use Disorder in the Past Year among People Aged 12 or Older: 2015-2019




+ Difference between this estimate and the 2019 estimate is statistically significant at the .05 level.

Figure 37. Alcohol Use Disorder in the Past Year among People Aged 12 or Older: 2002-2019




+ Difference between this estimate and the 2019 estimate is statistically significant at the .05 level.

Figure 39. Marijuana Use Disorder in the Past Year among People Aged 12 or Older: 2002-2019



+ Difference between this estimate and the 2019 estimate is statistically significant at the .05 level.

Figure 38. Illicit Drug Use Disorder in the Past Year among People Aged 12 or Older: 2015-2019



+ Difference between this estimate and the 2019 estimate is statistically significant at the .05 level.

Figure 1. Drug overdose death rates for adolescents aged 15–19, by sex: United States, 1999–2015






Figure 3. Age-adjusted drug overdose death rates involving opioids, by type of opioid: United States, 1999–2018




Positive Social Supports and Family Engagement



Rationale

- Family factors that reduce substance use in adolescents:
 - Warm and supportive family environment
 - Parental monitoring
- School factors that reduce substance use in adolescents:
 - Warm, supportive, nonpunitive school environment that promotes prosocial behavior
 - Schools that promote opportunities for positive socialization and interrupt deviant peer clustering
- Community factors that reduce substance use in adolescents:
 - Absence of substance-based advertising
 - Juvenile justice approaches that promote restorative practices, skill building, counseling and multiple coordinated services




Family Engagement

- Reduction of parent-child conflict
- Improved care-giver mental health
- Improved rates of treatment adherence and completion
- Longer duration of abstinence from substance use
- Fewer relapses




Kumpfer 2003; Liddle 2004; Steinglass 2009; Copello 2005; Smith 2004




Barriers to Family Engagement

- Family member ambivalence about need for treatment
 - Most youth in need of treatment, do not consider their substance use problematic
 - Perception that substance use is a normative behavior in adolescence
- Lack of understanding of what treatment entails




Barriers to Family Engagement

- Family member disengagement after initiation of treatment
 - Family members see substance use disorders as episodic, rather than as a chronic relapsing-remitting illness
- Perception that substance use is “the child’s problem/fault”
- Transportation and scheduling issues
- Care givers’ own addiction or mental illness




Family Engagement and SUD Treatment

- In a survey of adolescents with various mental health problems 91% cited that their **parent(s)** were the most influential to their decision to enter treatment (vs. teachers, counselors, doctors, etc).
- Parental involvement in adolescent treatment predicts improved treatment retention
- Greater parental utilization of treatment services is associated with decreased adolescent substance use



Family Engagement and SUD Treatment

- Differences in the subjective perception of external treatment pressures result in different therapeutic alliance outcomes
 - Parental threats → fear of consequences (**less internal motivation**) → less therapeutic alliance
 - Parental concern → acceptance that treatment is necessary for a better future (**more internal motivation**) → more therapeutic alliance



Provider Guidance on Family Engagement

- Educate families on neurodevelopmental issues related to substance use
- Provide definitions of substance use disorders (when is substance use problematic?)
- Educate families on SUD as a chronic, remitting-relapsing illness
- Counsel families about evidence-based treatment options for SUDs
- Provide guidance on externally motivated treatment
- Recommend local treatment programs that utilize evidence-based treatment strategies



Conclusions

- Problematic substance use among adolescents is often underrecognized by healthcare providers due to lack of education on this topic
- Mental illness and substance use are major contributors to health burden among youth and negatively impact outcomes in adulthood
- Rates of alcohol use and cigarette smoking among U.S. adolescents continue to decline
- Nicotine vaping and daily cannabis use are areas of concern
- While rates of Rx drug misuse are on the decline, rates of overdose death are rising among youth
- Positive social supports and family engagement are key factors for addressing problematic substance use in adolescents



Thank You!


Veronika Mesheriakova, MD

Assistant Professor

Pediatrics and Adolescent Medicine,
University of California, San Francisco

Director



UCSF Youth Outpatient Substance Use
Program (YoSUP)



In Our Last Few Moments...

- Follow-up email
- PowerPoint slides
- On-demand access 24/7
- Brief survey
- Certificate of Completion brief application (1 NAADAC CE)

sbirt.webs.com/adolescent-epidemiology





Related Products & Resources from the ATTC Network

- [CLAS Standards in Behavioral Health: Working with Youth and Adolescents](#) (Recorded webinar)
- [Understanding Latino Youth Recovery: Issues, Assets and Creating Resiliency](#) (Recorded webinar)
- Adolescent Brain Maturation and Health: Intersections on the Developmental Highway
 - [Recorded presentation](#)
 - [Handouts](#)
- [Effects on Marijuana Use on Developing Adolescents](#) (Recorded webinar)
- [Vaping Overview and CATCH My Breath Program](#) (Recorded webinar)
- [Vaping 2: Education vs Punishment Using Deferred Citation](#) (Recorded webinar)
- [Understanding Suicide Part 2 Adolescents and the Changing Brain](#) (Recorded webinar)

attcnetwork.org/centers/global-attc/tay-webinar-series



ATTC

Keep in Touch with the ATTC Network

- ATTC Network Office publishes the Messenger monthly
 - Subscribe: <https://attcnetwork.org/centers/global-attc/subscribe-attc-messenger>



@ATTcnetwork



@attcNetwork



ATTC

Addiction Technology Transfer Center Network
Funded by Substance Abuse and Mental Health Services Administration

the **Messenger** October 2020

October is Domestic Violence Awareness Month

In Honor of Domestic Violence Awareness Month:

Responding to Substance Use Coercion in

Treatment and Recovery Services

By Carole Warshaw, MD, and Gabriela Zapata-Alma,
LCSW, CADC,

Intimate Partner Violence (IPV) Can Have Profound

include deliberately introducing a partner to substances, forcing or coercing a partner to use, interfering with treatment, controlling medication; sabotaging recovery efforts; threatening a partner with withdrawal, and leveraging the stigma associated with substance use to discredit a partner with potential sources of safety and support.

attcnetwork.org/centers/global-attc/tay-webinar-series



Related Products & Resources from the PTTC Network

- [Underage Alcohol Use: An Overview of Data and Strategies](#) (Recorded webinar)
- [Youth Opioid Addiction: What Preventionists Need to Know](#) (Recorded webinar)
- [Selecting and Implementing Evidence-Based Practices to Address Substance Misuse Among Young Adults: Webinar on SAMHSA's Resource Guide](#)
- [Preventing Youth Vaping \(Webinar Series\) Part 1 of 2: The Extent and Risk Factors for Youth Vaping](#) (Recorded webinar)
- [Preventing Youth Vaping Part 2 of 2: Policy Recommendations and Promising Practices for Addressing Youth Vaping](#) (Recorded webinar)
- [The Benefits of Engaging Youth in Communities: Insights and Evidence from Developmental Science](#) (Recorded webinar)
- [Vaping and LGBTQ Youth](#) (Recorded webinar)
- [Informing Prevention 6-Part Webinar Series on Adolescents: Mountain Plains PTTC](#)
- [Adolescent SBIRT Pocket Card](#)

attcnetwork.org/centers/global-attc/tay-webinar-series



Keep In Touch with the PTTC Network

- PTTC Network Office publishes the *PTTC POST* monthly
 - Please Subscribe:
<https://pttcnetwork.org/centers/global-pttc/pttc-subscription-page>



@PTTCnetwork



@PreventionTTCnetwork



pttcnetwork.org/centers/global-pttc/tay-webinar-series



TRANSITIONAL AGE YOUTH (TAY)
webinar series

Who's Doing What?: The Epidemiology of Adolescent Substance Use

Produced in Partnership by:



Network Coordinating Office

ATTC

Addiction Technology Transfer Center Network
Funded by Substance Abuse and Mental Health Services Administration



Interdisciplinary Leaders in
Substance Use Education,
Research, Care and Policy



by



attnetwork.org/centers/global-atcc/tay-webinar-series