

Adult ASAM Placement Criteria Crosswalk

Adult ASAM Worksheet	Level 0.5: Early Intervention	OTP Level 1: Opioid Treatment Program	Level 1: Outpatient Services	Level 2.1: Intensive Outpatient Treatment	Level 2.5: Partial Hospitalization	Level 3.1: Clinically – Managed Low-Intensity Residential Services	Level 3.3: Clinically-Managed Population Specific High-Intensity Residential Services	Level 3.5: Clinically Managed High Intensity Residential Services	Level 3.7: Medically Monitored Intensive Inpatient Services	Level 4: Medically Managed Intensive Inpatient Services
Dimension 1: Acute Intoxication and/or Withdrawal Potential	No withdrawal risk	Physiologically dependent on opioids and requires OTP to prevent withdrawal	Manageable at Level 1-WM	Manageable at Level 2-WM	Manageable at Level 2-WM	Concurrently receiving Level 1-WM or Level 2-WM services	If withdrawal is present, manageable at Level 3.2-WM.	If withdrawal is present, manageable at Level 3.2-WM.	At high risk of withdrawal, but manageable at Level 3.7-WM and does not require the full resources of a licensed hospital	At high risk of withdrawal and requires Level 4-WM and the full resources of a licensed hospital
Dimension 2: Biomedical Conditions and Complications	None or very stable	None or manageable with outpatient medical monitoring	None or very stable, or is receiving concurrent medical monitoring	None or not a distraction from treatment	None or not sufficient to distract from treatment	None or stable, or receiving concurrent medical monitoring	None or stable, or receiving concurrent medical monitoring	None or stable, or receiving concurrent medical monitoring	Requires 24 hours medical monitoring but not intensive treatment	Requires 24 hour medical and nursing care and the full resources of a licensed hospital
Dimension 3: Emotional, Behavioral or Cognitive Conditions and Complications	None or very stable	None or manageable in an outpatient structured environment	None or very stable, or is receiving concurrent medical monitoring	Mild severity with the potential to distract from recovery; needs monitoring	Mild to Moderate severity with the potential to distract from recovery; needs stabilization	None or minimal; not distracting to recovery. If stable, a co-occurring capable program is appropriate. If not, a co-occurring enhanced program is required	Mild to moderate severity, needs structure to focus on recovery. Tx should be designed to address significant cognitive deficits. If stable, a co-occurring program is appropriate. If not, a co-occurring enhanced program is required	Demonstrates repeated inability to control impulses, or unstable and dangerous signs/sx require stabilization. Other functional deficits need stabilization and 24-hour setting to prepare for community integration and continuing care. A co-occurring enhanced setting required for those with severe and chronic MI	Moderate severity; needs a 24-hour structured setting, if the consumer has a co-occurring mental disorder, requires concurrent MH services in a medically monitored setting	Because of severe and unstable problems; requires 24 hour psychiatric care with concomitant addiction tx (co-occurring enhanced)
Dimension 4: Readiness to Change	Consumer is willing to explore how current alcohol, tobacco, other drug or medication use and/or high risk behaviors may affect personal goals	Ready to change the negative effects of opioid use, but is not ready for total abstinence from illicit prescription or non-prescription drug use	Ready for recovery but needs motivating and monitoring strategies to strengthen readiness, or needs ongoing monitoring and disease management . Or high severity in this dimension but not in other dimensions. Needs Level 1 motivational enhancement strategies.	Has variable engagement in treatment, ambivalence, or a lack of awareness of the substance use or MH problem, and requires a structured program several times a week to promote progress through the stages of change	Has poor engagement in tx, significant ambivalence or a lack of awareness of the substance use or MH problem, requiring a near-daily structured program or intensive engagement service to promote progress through the stages of change	Open to recovery, but needs a structured environment to maintain therapeutic gains	Has little awareness and needs interventions available only at Level 3.3 to engage and stay in tx. If there is high severity in Dimension 4 but not in any other dimension, motivational enhancement strategies should be provided in Level 1	Has marked difficulty with, or oppositional to, tx, with dangerous consequences. If there is high severity in Dimension 4 but not in any other dimension, motivation enhancement strategies should be provided in Level 1	Low interest in tx and impulse control is poor, despite negative consequences; needs motivating strategies only safely available in a 24-hour structured setting. If there is high severity in Dimension 4 but not in any other dimension, motivational enhancement strategies should be provided in Level 1	Problems in this dimension do not qualify the consumer for Level 4 services. If the consumer's only severity is in Dimension 4,5 and/or 6 without high severity in Dimensions 1,2, and /or 3, then the consumer does not qualify for Level 4

Dimensions 5 and 6 on next page

Adult ASAM Placement Criteria Crosswalk

<p>Dimension 5: Relapse, Continued Use or Continued Problem Potential</p>	<p>Needs an understanding of, or skills to change, current alcohol, tobacco, other drug, or medication use patterns and/or high risk behavior</p>	<p>At high risk of relapse or continued use without OTP and structured therapy to promote tx progress</p>	<p>Able to maintain abstinence or control use and/or addictive behaviors and pursue recovery or motivational goals with minimal support</p>	<p>Intensification of addiction or MH sx indicate a high likelihood of relapse or continued use or without close monitoring and support several times a week</p>	<p>Intensification of addiction or MH sx, despite active participation in a Level 1 or 2.1 program; indicates a high likelihood of relapse or continued use or continued problems without near daily monitoring and support</p>	<p>Understands relapse, but needs structure to maintain therapeutic gains</p>	<p>Has little awareness and needs interventions available only at Level 3.3 to prevent continued use, with imminent dangerous consequences, because of cognitive deficits or comparable dysfunction</p>	<p>Has no recognition of the skills needed to prevent continued use, with imminent dangerous consequences</p>	<p>Unable to control use, with imminently dangerous consequences despite active participation at less intensive levels of care</p>	<p>Problems in this dimension do not qualify the consumer for Level 4 services. See further explanation in Dimension 4</p>
<p>Dimension 6: Recovery Environment</p>	<p>Social support system or significant others increase the risk of personal conflict about alcohol, tobacco, and/or other drug use</p>	<p>Recovery environment is supportive and/or consumer has skills to cope</p>	<p>Recovery environment is supportive and/or the consumer has skills to cope</p>	<p>Recovery environment is not supportive, but with structure and support the consumer can cope</p>	<p>Recovery environment is not supportive, but with structure and support and relief from the home environment, the consumer can cope</p>	<p>Environment is dangerous, but recovery is achievable if Level 3.1, 24-hour structure is available</p>	<p>Environment is dangerous and consumer needs 24-hour structure to learn to cope</p>	<p>Environment is dangerous and the consumer lacks skills to cope outside of a highly structured 24-hour setting</p>	<p>Environment is dangerous and the consumer lacks skills to cope outside of a highly structured 24-hour setting</p>	<p>Problems in this dimension do not qualify the consumer for Level 4 services. See further explanation in Dimension 4</p>

To select the correct level of care, choose the highest level of care that has two or more criteria met. Review admission criteria for chosen level of care to ensure proper clinical fit.

NAME or SM ID#: _____

CURRENT ASAM LEVEL: _____

RECOMMENDED LEVEL OF CARE AND SUBSTANCE ABUSE SERVICES: _____

CLINICIAN SIGNATURE: _____

DATE: _____